

Bianchi

Scaffidi, P., T. Misteli, and M.E. Bianchi.
Release of chromatin protein HMGB1 by necrotic cells triggers inflammation.
Nature 418: 191-195 (2002).

Gardella, S., C. Andrei, D. Ferrera, L.V. Lotti, M.R. Torrisi, M.E. Bianchi, and A. Rubartelli. The nuclear protein HMGB1 is secreted by monocytes via a non-classical, vesicle-mediated secretory pathway.
EMBO Rep. 3: 995-1001 (2002).

Agresti, A., and M.E. Bianchi.
HMGB proteins and gene expression.
Curr. Opin. Genet. Dev. 13: 170-178 (2003).

Agresti, A., R. Lupo, M.E. Bianchi, and S. Muller.
HMGB1 interacts differentially with members of the Rel family of transcription factors. Biochem. Biophys. Res. Commun. 302:421-426 (2003).

Bonaldi, T., F. Talamo, P. Scaffidi, D. Ferrera, A. Porto, A. Bachi, A. Rubartelli, A. Agresti, and M.E. Bianchi.
Monocytic cells hyperacetylate chromatin protein HMGB1 to redirect it towards secretion. EMBO J. 22: 5551-5560 (2003).

Guazzi, S., A. Strangio, A.T. Franzi, and M.E. Bianchi.
HMGB1, an architectural chromatin protein and extracellular signalling factor, has a spatially and temporally restricted expression pattern in mouse brain.
Gene. Expr. Patterns 3: 29-33 (2003).

Krynetski, E.Y., N.F. Krynetskaia, M.E. Bianchi, and W.E. Evans.
A nuclear protein complex containing high mobility group proteins B1 and B2, heat shock cognate protein 70, ERp60, and glyceraldehyde-3-phosphate dehydrogenase is involved in the cytotoxic response to DNA modified by incorporation of anticancer nucleoside analogues. Cancer Res. 63:100-106 (2003).

Lalli, E., K. Ohe, E. Latorre, M.E. Bianchi, and P. Sassone-Corsi.
Sexy splicing: regulatory interplays governing sex determination from Drosophila to mammals.
J. Cell Sci. 116:441-445 (2003).

Pallier, C., P. Scaffidi, S. Chopineau-Proust, A. Agresti, P. Nordmann, M.E. Bianchi, and V. Marechal.
Association of chromatin proteins high mobility group box (HMGB) 1 and HMGB2 with mitotic chromosomes.
Mol. Biol. Cell. 14: 3414-3426 (2003).

Palumbo, R., and M.E. Bianchi.
High mobility group box 1 protein, a cue for stem cell recruitment.
Biochem. Pharmacol. 68: 1165-1170 (2004).

Ronfani, L., and M.E. Bianchi.

Molecular mechanisms in male determination and germ cell differentiation.
Cell. Mol. Life Sci. 61: 1907-1925 (2004).

Rovere-Querini, P., A. Capobianco, P. Scaffidi, B. Valentinis, F. Catalanotti, M. Giazzon, I.E. Dumitriu, S. Muller, M. Iannacone, C. Traversari, M.E. Bianchi, and A.A. Manfredi.

HMGB1 is an endogenous immune adjuvant released by necrotic cells.
EMBO Rep. 5: 825-830 (2004).

Sawchuk, D.J., J. Mansilla-Soto, C. Alarcon, N.C. Singha, H. Langen, M.E. Bianchi, S.P. Lees-Miller, M.C. Nussenzweig, and P. Cortes.
Ku70/Ku80 and DNA-dependent protein kinase catalytic subunit modulate RAG-mediated cleavage: implications for the enforcement of the 12/23 rule.
J. Biol. Chem. 279: 29821-29831. (2004).

Bianchi, M.E.

Significant (re)location: how to use chromatin and/or abundant proteins as messages of life and death.
Trends Cell Biol. 14: 287-293 (2004).

Bianchi, M.E., and A. Manfredi.

Chromatin and cell death.
Biochim. Biophys. Acta 1677: 181-186 (2004).

Knapp, S., S. Muller, G. Digilio, T. Bonaldi, M.E. Bianchi, and G. Musco.
The long acidic tail of high mobility group box 1 (HMGB1) protein forms an extended and flexible structure that interacts with specific residues within and between the HMG boxes. Biochemistry. 43: 11992-11997 (2004).

Muller, S., L. Ronfani, and M.E. Bianchi.

Regulated expression and subcellular localization of HMGB1, a chromatin protein with a cytokine function.
J. Intern. Med. 255: 332-343 (2004).

Palumbo, R., M. Sampaolesi, F. De Marchis, R. Tonlorenzi, S. Colombetti, A. Mondino, G. Cossu, and M.E. Bianchi.

Extracellular HMGB1, a signal of tissue damage, induces mesoangioblast migration and proliferation.
J. Cell Biol. 164: 441-449 (2004).

Agresti, A., P. Scaffidi, A. Riva, V.R. Caiolfa, and M.E. Bianchi.

GR and HMGB1 interact only within chromatin and influence each other's residence time. Mol. Cell 18:109-121 (2005).

Andrassy, M., H.C. Volz, J.C. Igwe, B. Funke, S.N. Eichberger, Z. Kaya, S. Buss, F. Bianchi, M.E., and A. Agresti.

HMG proteins: dynamic players in gene regulation and differentiation.

Curr. Opin Genet Dev. 15:496-506 (2005).

Bianchi, M.E., and A. Agresti.

HMG proteins: dynamic players in gene regulation and differentiation.

Curr Opin. Genet. Dev. 15: 496-506 (2005).

Dumitriu, I.E., P. Baruah, M.E. Bianchi, A.A. Manfredi, and P. Rovere-Querini
Requirement of HMGB1 and RAGE for the maturation of human plasmacytoid
dendritic cells.

Eur. J. Immunol. 35: 2184-2190 (2005).

Dumitriu, I.E., P. Baruah, A.A. Manfredi, M.E. Bianchi, and P. Rovere-Querini.
HMGB1: guiding immunity from within.

Trends Immunol. 26: 381-387 (2005).

Dumitriu, I.E., P. Baruah, B. Valentinis, R.E. Voll, M. Herrmann, P.P. Nawroth,
B. Arnold, M.E. Bianchi, A.A. Manfredi, and P. Rovere-Querini.

Release of high mobility group box 1 by dendritic cells controls T cell
activation via the receptor for advanced glycation end products.

J. Immunol. 174: 7506-7515 (2005).

Giavara, S., E. Kosmidou, M.P. Hande, M.E. Bianchi, A. Morgan, F. d'Adda di
Fagagna, and S.P. Jackson.

Yeast Nhp6A/B and mammalian Hmgb1 facilitate the maintenance of genome
stability.

Curr. Biol. 15: 68-72 (2005).

Limana, F., A. Germani, A. Zacheo, J. Kajstura, A. Di Carlo, G. Borsellino, O.
Leoni, R. Palumbo, L. Battistini, R. Rastaldo, S. Muller, G. Pompilio, P.

Anversa, M.E. Bianchi, and M.C. Capogrossi.

Exogenous high-mobility group box 1 protein induces myocardial
regeneration after infarction via enhanced cardiac C-kit⁺ cell proliferation and
differentiation.

Circ. Res. 97:e73-83 (2005).

Porto, A., R. Palumbo, M. Pieroni, G. Aprigliano, R. Chiesa, F. Sanvito, A.
Maseri, and M.E. Bianchi.

Smooth muscle cells in human atherosclerotic plaques secrete and
proliferate in response to high mobility group box 1 protein.

FASEB J. 20: 2565-2566 (2006).

Bosisio, D., I. Marazzi, A. Agresti, N. Shimizu, M.E. Bianchi, and G. Natoli.

A hyper-dynamic equilibrium between promoter-bound and nucleoplasmic
dimers controls NF-kappaB-dependent gene activity.

EMBO J. 25: 798-810 (2006).

Bianchi, M.E.

DAMPs, PAMPs and alarmins: all we need to know about danger.

J. Leukoc. Biol. 81: 1-5 (2007).

Bianchi, M.E., and A.A. Manfredi.

High-mobility group box 1 (HMGB1) protein at the crossroads between innate and adaptive immunity.

Immunol. Rev. 220: 35-46 (2007).

Chavakis, E., A. Hain, M. Vinci, G. Carmona, M.E. Bianchi, P. Vajkoczy, A.M. Zeiher, T. Chavakis, and S. Dimmeler.

High-mobility group box 1 activates integrin-dependent homing of endothelial progenitor cells.

Circ. Res. 100: 204-212 (2007).

De Mori, R., S. Straino, A. Di Carlo, A. Mangoni, G. Pompilio, R. Palumbo, M.E. Bianchi, M.C. Capogrossi, and A. Germani.

Multiple effects of high mobility group box protein 1 in skeletal muscle regeneration. Arterioscler. Thromb. Vasc. Biol. 27: 2377-2383 (2007).

Dumitriu, I.E., M.E. Bianchi, M. Bacci, A.A. Manfredi, and P. Rovere-Querini. The secretion of HMGB1 is required for the migration of maturing dendritic cells.

J. Leukoc. Biol. 81: 84-91 (2007).

Faraco, G., S. Fossati, M.E. Bianchi, M. Patrone, M. Pedrazzi, B. Sparatore, F. Moroni, and A. Chiarugi..

High mobility group box 1 protein is released by neural cells upon different stresses and worsens ischemic neurodegeneration in vitro and in vivo.

J. Neurochem. 103: 590-603 (2007).

Ivanov, S., A.M. Dragoi, X. Wang, C. Dallacosta, J. Louten, G. Musco, G. Sitia, G.S. Yap, Y. Wan, C.A. Biron, M.E. Bianchi, H. Wang, and W.M. Chu.

A novel role for HMGB1 in TLR9-mediated inflammatory responses to CpG-DNA.

Blood 110: 1970-1981 (2007).

Mollica, L., F. De Marchis, A. Spitaleri, C. Dallacosta, D. Pennacchini, M.

Zamai, A. Agresti, L. Trisciuglio, G. Musco, and M.E. Bianchi..

Glycyrrhizin binds to high-mobility group box 1 protein and inhibits its cytokine activities.

Chem. Biol. 14: 431-441 (2007).

Orlova, V.V., E.Y. Choi, C. Xie, E. Chavakis, A. Bierhaus, E. Ihanus, C.M.

Ballantyne, C.G. Gahmberg, M.E. Bianchi, P.P. Nawroth, and T. Chavakis.

A novel pathway of HMGB1-mediated inflammatory cell recruitment that requires Mac-1-integrin.

EMBO J. 26: 1129-1139 (2007).

Palumbo, R., B.G. Galvez, T. Pusterla, F. De Marchis, G. Cossu, K.B. Marcu, and M.E. Bianchi.

Cells migrating to sites of tissue damage in response to the danger signal HMGB1 require NF-kappaB activation.

J. Cell Biol. 179: 33-40 (2007).

Raucci, A., R. Palumbo, and M.E. Bianchi.
HMGB1: a signal of necrosis.
Autoimmunity 40: 285-289 (2007).

Sitia, G., M. Iannacone, S. Muller, M.E. Bianchi, and L.G. Guidotti.
Treatment with HMGB1 inhibitors diminishes CTL-induced liver disease in
HBV transgenic mice.
J. Leukoc. Biol. 81: 100-107 (2007).

Taniguchi, N., K. Yoshida, T. Ito, M. Tsuda, Y. Mishima, T. Furumatsu, L.
Ronfani, K. Abeyama, K. Kawahara, S. Komiya, I. Maruyama, M. Lotz, M.E.
Bianchi, and H. Asahara.
Stage-specific secretion of HMGB1 in cartilage regulates endochondral
ossification.
Mol. Cell Biol. 27: 5650-5663 (2007).

Sessa, L., and M.E. Bianchi..
The evolution of High Mobility Group Box (HMGB) chromatin proteins in
multicellular animals.
Gene 387: 133-140 (2007).

Raucci, A., S. Cugusi, A. Antonelli, S.M. Barabino, L. Monti, A. Bierhaus, K.
Reiss, P. Saftig, and M.E. Bianchi.
A soluble form of the receptor for advanced glycation endproducts (RAGE) is
produced by proteolytic cleavage of the membrane-bound form by the
shedase a disintegrin and metalloprotease 10 (ADAM10).
FASEB J. 22: 3716-3727 (2008).

Manfredi, A.A., A. Capobianco, A. Esposito, F. De Cobelli, T. Canu, A.
Monno, A. Raucci, F. Sanvito, C. Doglioni, P.P. Nawroth, A. Bierhaus, M.E.
Bianchi, P. Rovere-Querini, and A. Del Maschio.
Maturing dendritic cells depend on RAGE for in vivo homing to lymph nodes.
J. Immunol. 180: 2270-2275 (2008).

Autschbach, S.T. Pleger, I.K. Lukic, F. Bea, S.E. Hardt, P.M. Humpert, M.E.
Bianchi, H. Mairbaur, P.P. Nawroth, A. Remppis, H.A. Katus, and A.
Bierhaus.
High-mobility group box-1 in ischemia-reperfusion injury of the heart.
Circulation 117: 3216-3226 (2008).

Urbonaviciute, V., B.G. Furnrohr, S. Meister, L. Munoz, P. Heyder, F. De
Marchis, M.E. Bianchi, C. Kirschning, H. Wagner, A.A. Manfredi, J.R. Kalden,
G. Schett, P. Rovere-Querini, M. Herrmann, and R.E. Voll.
Induction of inflammatory and immune responses by HMGB1-nucleosome
complexes: implications for the pathogenesis of SLE.
J. Exp. Med. 205: 3007-3018. (2008).

Straino, S., A. Di Carlo, A. Mangoni, R. De Mori, L. Guerra, R. Maurelli, L. Panacchia, F. Di Giacomo, R. Palumbo, C. Di Campli, L. Uccioli, P. Biglioli, M.E. Bianchi, M.C. Capogrossi, and A. Germani.

High-mobility group box 1 protein in human and murine skin: involvement in wound healing.

J. Invest. Dermatol. 128: 1545-1553 (2008).

Bianchi, M.E.

HMGB1 loves company.

J. Leukoc. Biol. (2009).

Bianchi, M.E., and A.A. Manfredi..

Immunology. Dangers in and out.

Science. 323: 1683-1684 (2009).

Campana, L., L. Bosurgi, M.E. Bianchi, A.A. Manfredi, and P. Rovere-Querini. Requirement of HMGB1 for stromal cell-derived factor-1/CXCL12-dependent migration of macrophages and dendritic cells.

J. Leukoc. Biol. (2009).

Cassetta, L., O. Fortunato, L. Adduce, C. Rizzi, J. Hering, P. Rovere-Querini, M.E. Bianchi, M. Alfano, and G. Poli.

Extracellular high mobility group box-1 inhibits R5 and X4 HIV-1 strains replication in mononuclear phagocytes without induction of chemokines and cytokines.

Aids. 23: 567-577 (2009).

Lolmede, K., L. Campana, M. Vezzoli, L. Bosurgi, R. Tonlorenzi, E. Clementi, M.E. Bianchi, G. Cossu, A.A. Manfredi, S. Brunelli, and P. Rovere-Querini.

Inflammatory and alternatively activated human macrophages attract vessel-associated stem cells, relying on separate HMGB1- and MMP-9-dependent pathways.

J. Leukoc. Biol. 85: 779-787 (2009).

Manfredi, A.A., A. Capobianco, M.E. Bianchi, and P. Rovere-Querini.

Regulation of dendritic- and T-cell fate by injury-associated endogenous signals.

Crit. Rev. Immunol. 29: 69-86 (2009).

Palumbo, R., F. De Marchis, T. Pusterla, A. Conti, M. Alessio, and M.E. Bianchi.

Src family kinases are necessary for cell migration induced by extracellular HMGB1.

J. Leukoc. Biol. (2009).

Taniguchi, N., B. Carames, L. Ronfani, U. Ulmer, S. Komiya, M.E. Bianchi, and M. Lotz. Aging-related loss of the chromatin protein HMGB2 in articular cartilage is linked to reduced cellularity and osteoarthritis.

Proc. Natl. Acad. Sci. U. S. A. 106: 1181-1186 (2009).