

SITIA

Fassio A, Sitia R.

Formation, isomerisation and reduction of disulphide bonds during protein quality control in the endoplasmic reticulum.

Histochem Cell Biol. 117:151-157 (2002).

Anelli T, Alessio M, Mezghrani A, Simmen T, Talamo F, Bachi A, Sitia R.
ERp44, a novel endoplasmic reticulum folding assistant of the thioredoxin family.

EMBO J. 21: 835-844 (2002).

Angelini G, Gardella S, Ardy M, Ciriolo MR, Filomeni G, Di Trapani G, Clarke F, Sitia R, Rubartelli A.

Antigen-presenting dendritic cells provide the reducing extracellular microenvironment required for T lymphocyte activation.

Proc. Natl. Acad. Sci. U. S. A. 99:1491-1496 (2002).

* Sitia R, Braakman I.

Quality control in the endoplasmic reticulum protein factory.

Nature 426: 891-894 (2003).

* Anelli T, Alessio M, Bachi A, Bergamelli L, Bertoli G, Camerini S, Mezghrani A, Ruffato E, Simmen T, Sitia R.

Thiol-mediated protein retention in the endoplasmic reticulum: the role of ERp44.

EMBO J. 22: 5015-5022 (2003).

Tagliavacca L, Anelli T, Fagioli C, Mezghrani A, Ruffato E, Sitia R.

The making of a professional secretory cell: architectural and functional changes in the ER during B lymphocyte plasma cell differentiation.

J. Biol. Chem. 384:1273-1277 (2003).

* van Anken E, Romijn EP, Maggioni C, Mezghrani A, Sitia R, Braakman I, Heck AJ.

Sequential waves of functionally related proteins are expressed when B cells prepare for antibody secretion.

Immunity 18:243-253 (2003).

* Cunnea PM, Miranda-Vizuete A, Bertoli G, Simmen T, Damdimopoulos AE, Hermann S, Leinonen S, Huikko MP, Gustafsson JA, Sitia R, Spyrou G.

ERdj5, an endoplasmic reticulum (ER)-resident protein containing DnaJ and thioredoxin domains, is expressed in secretory cells or following ER stress.

J. Biol. Chem. 278: 1059-1066 (2003).

* Sitia R, Molteni SN.

Stress, protein (mis)folding, and signaling: the redox connection.
Sci. STKE. 2004: pe27 (2004).

* Molteni SN, Fassio A, Ciriolo MR, Filomeni G, Pasqualetto E, Fagioli C, Sitia R.

Glutathione limits Ero1-dependent oxidation in the endoplasmic reticulum.
J. Biol. Chem. 279: 32667-32673 (2004).

* Bertoli G, Simmen T, Anelli T, Molteni SN, Fesce R, Sitia R.

Two conserved cysteine triads in human Ero1alpha cooperate for efficient disulfide bond formation in the endoplasmic reticulum.

J. Biol. Chem. 279: 30047-30052 (2004).

Piccini A, Fassio A, Pasqualetto E, Vitali A, Borghi R, Palmieri D, Nacmias B, Sorbi S, Sitia R, Tabaton M.

Fibroblasts from FAD-linked presenilin 1 mutations display a normal unfolded protein response but overproduce Abeta42 in response to tunicamycin.

Neurobiol. Dis. 15: 380-386 (2004).

* Molinari M, Sitia R.

The secretory capacity of a cell depends on the efficiency of endoplasmic reticulum-associated degradation.

Curr. Top. Microbiol. Immunol. 300: 1-15 (2005).

* Orsi A, Fioriti L, Chiesa R, Sitia R.

Conditions of endoplasmic reticulum stress favor the accumulation of cytosolic prion protein.

J. Biol. Chem. 281:30431-30438 (2006).

* Mattioli L, Anelli T, Fagioli C, Tacchetti C, Sitia R, Valetti C.

ER storage diseases: a role for ERGIC-53 in controlling the formation and shape of Russell bodies.

J. Cell. Sci. 119:2532-2541 (2006).

* Otsu M, Bertoli G, Fagioli C, Guerini-Rocco E, Nerini-Molteni S, Ruffato E, Sitia R.

Dynamic retention of Ero1alpha and Ero1beta in the endoplasmic reticulum by interactions with PDI and ERp44.

Antioxid. Redox. Signal. 8: 274-282 (2006).

Benham AM, Sitia R.

The diversity of oxidative protein folding.

Antioxid. Redox. Signal. 8: 271-273 (2006).

* Cenci S, Mezghrani A, Cascio P, Bianchi G, Cerruti F, Fra A, Lelouard H, Masciarelli S, Mattioli L, Oliva L, Orsi A, Pasqualetto E, Pierre P, Ruffato

E, Tagliavacca L, Sitia R.
Progressively impaired proteasomal capacity during terminal plasma cell differentiation.
EMBO J. 25: 1104-1113 (2006).

* Sitia R, Palladini G, Merlini G.
Bortezomib in the treatment of AL amyloidosis: targeted therapy?
Haematologica. 92: 1302-1307 (2007).

*Orsi A, Sitia R.
Interplays between covalent modifications in the endoplasmic reticulum increase conformational diversity in nascent prion protein.
Prion 1: 236-42 (2007).

Anelli T, Ceppi S, Bergamelli L, Cortini M, Masciarelli S, Valetti C, Sitia R.
Sequential steps and checkpoints in the early exocytic compartment during secretory IgM biogenesis.
EMBO J. 26:4177-4188 (2007).

Otsu M, Sitia R.
Diseases originating from altered protein quality control in the endoplasmic reticulum.
Curr. Med. Chem. 14: 1639-1652 (2007).

Bánhegyi G, Baumeister P, Benedetti A, Dong D, Fu Y, Lee AS, Li J, Mao C, Margittai E, Ni M, Paschen W, Piccirella S, Senesi S, Sitia R, Wang M, Yang W.
Endoplasmic reticulum stress.
Ann. N. Y. Acad. Sci. 1113:58-71 (2007).

* Cenci S, Sitia R.
Managing and exploiting stress in the antibody factory.
FEBS Lett. 2007 Jul;581:3652-7

Cenci S, Pengo N, Sitia R.
Proteotoxic stress and cell lifespan control.
Mol. Cell. 26: 323-3288 (2008).

Wang L, Wang L, Vavassori S, Li S, Ke H, Anelli T, Degano M, Ronzoni R, Sitia R, Sun F, Wang CC Crystal structure of human ERp44 shows a dynamic functional modulation by its carboxy-terminal tail.
EMBO Rep. 9: 642-627 (2008).

Fraldi A, Zito E, Annunziata F, Lombardi A, Cozzolino M, Monti M, Spanpanato C, Ballabio A, Pucci P, Sitia R, Cosma MP.
Multistep, sequential control of the trafficking and function of the multiple sulfatase deficiency gene product, SUMF1 by PDI, ERGIC-53 and ERp44.
Hum. Mol. Genet. 17: 2610-2621 (2008).

Nerini-Molteni S, Ferrarini M, Cozza S, Caligaris-Cappio F, Sitia R.
Redox homeostasis modulates the sensitivity of myeloma cells to
bortezomib.

Br. J. Haematol. 141: 494-503 (2008).

Cattaneo M, Otsu M, Fagioli C, Martino S, Lotti LV, Sitia R, Biunno I.
SEL1L and HRD1 are involved in the degradation of unassembled
secretory Ig-mu chains.

J. Cell. Physiol. 215: 794-802 (2008).

Cascio P, Oliva L, Cerruti F, Mariani E, Pasqualetto E, Cenci S, Sitia R.
Dampening Ab responses using proteasome inhibitors following in vivo B
cell activation.

Eur. J. Immunol. 38:658-667 (2008).

Masciarelli S, Sitia R.

Building and operating an antibody factory: redox control during B to
plasma cell terminal differentiation.

Biochim. Biophys. Acta. 1783: 578-588 (2008).

Anelli T, Sitia R.

Protein quality control in the early secretory pathway.

EMBO J. 27:315-327 (2008).

Rubartelli A, Sitia R.

Stress as an Intercellular Signal: The Emergence of Stress-Associated
Molecular Patterns (SAMP)

Antioxid. Redox. Signal. 2009 Mar.

Bianchi G, Oliva L, Cascio P, Pengo N, Fontana F, Cerruti F, Orsi A,
Pasqualetto E, Mezghrani A, Calbi V, Palladini G, Giuliani N, Anderson
KC, Sitia R, Cenci S.

The proteasome load versus capacity balance determines apoptotic
sensitivity of multiple myeloma cells to proteasome inhibition

Blood 113: 3040-9 (2009).